

## The prediction of athlete resting metabolic rate – is it time to reassess the method?

MacKenzie, Kristen

*Licence:*  
CC BY-NC-ND

[Link to output in Bond University research repository.](#)

*Recommended citation(APA):*  
MacKenzie, K. (2016). *The prediction of athlete resting metabolic rate – is it time to reassess the method?*.  
Poster session presented at The 17th International Congress of Dietetics Granada 2016, Granada, Spain.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

For more information, or if you believe that this document breaches copyright, please contact the Bond University research repository coordinator.



17th International Congress of Dietetics

GRANADA SPAIN 2016

7, 8, 9 and 10 September

www.icdgranada2016.com

Reference:

P - 010

**Title:** The prediction of athlete resting metabolic rate – is it time to reassess the method?

**Authors:** MacKenzie-Shalders, K<sup>1</sup>. McDonald, A<sup>2</sup>.

**Workcenter:** <sup>1</sup>Bond University, Gold Coast, AUSTRALIA.  
<sup>2</sup>Queensland University of Technology, Brisbane, AUSTRALIA.

**Introduction:** Effective energy prescription requires an accurate assessment of the athletes’ RMR. The use of published prediction equations using total body mass (TBM) or fat-free mass (FFM) with other covariates is common; but there is little evidence to validate their use or to determine which are most predictive in athlete groups.



**Methods:** This study compared measured resting metabolic rate (RMR) using indirect calorimetry to RMR using 17 prediction equations.

- Anthropometric and metabolic data was collected for 23 male rugby athletes
- A literature review was conducted for evidence relating to the measurement and prediction of RMR in athlete populations.
- Paired samples t-tests and root mean square prediction error (RMSPE) were used to compare measured and predicted RMR.




**Results:** The prediction equations significantly and systematically underestimated RMR in rugby players for all equations (p≤0.001).

- The Harris Benedict equation provided the most accurate estimate of RMR and predicted energy requirements within ± 189kcal/d (RMSPE).
- The commonly-recommended Cunningham equation using FFM was predictive ± 217 kcal/d (RMSPE).

	Mean kcal/d	Paired t-test		Mean Diff. kcal/d	RMSPE kcal/d
Measured RMR	2356 ±247	t	p-value		
Harris Benedict	2203 ±207	6.5	0.000	-154 ±112	189
Schofield	2189 ±204	7.1	0.000	-168 ±114	201
Cunningham	2187 ±154	5.9	0.000	-169 ±138	217

**Conclusions, discussion and/or practical application:**

- There are several sources of error that need to be addressed when applying these prediction equations to athletes.
- There is a need to identify the unique characteristics of athletes that act as covariates to develop effective prediction equations for athletes.



Queensland University of Technology



BOND UNIVERSITY

Cunningham, J. J. (1991). Body composition as a determinant of energy expenditure: a synthetic review and a proposed general prediction equation. *American Journal of Clinical Nutrition*, 54, 963 - 969.

Schofield, W. (1985). Predicting basal metabolic rate, new standard and review of previous work. *Human Nutrition, Clinical Nutrition*, 39c(Suppl 1), 5- 41.

Harris, J., & Benedict, F. (1919). A Biometric Study of Basal Metabolism in Man (pp. 1 - 266). Washington, DC. Carnegie: Institute of Washington.

Compher, C., Frankenfield, D., Keim, N., & Roth-Yousey, L. (2006). Best Practice Methods to Apply to Measurement of Resting Metabolic Rate in Adults: A Systematic Review. *Journal of the American Dietetic Association*, 106, 881-903.